arXiv:hep-th/9511033v1 6 Nov 1995

KUNS-1368
HE(TH) 95/18
hep-th/9511033
July, 1994

The Universal Renormalization Factors Z_1/Z_3 and Color Confinement Condition in Non-Abelian Gauge Theory*

Taichiro KUGO†

Department of Physics, Kyoto University, Kyoto 606-01, Japan

Abstract

The ratio Z_1/Z_3 of vertex and wave-function renormalization factors, which is universal (*i.e.*, matter-independent), is shown to equal $1+u$ which gives the residue of the scalar pole $\propto p_\mu p_\nu/p^2$ of 2-point function $\langle\, D_\mu c\; D_\nu \bar{c}\,\rangle$. This relation is interesting since $1+u=0$ has been known to give a sufficient condition for color confinement. We also give an argument that, when $1+u=0$ holds, it will be realized by the disappearance of the massless gauge boson pole and is related with the restoration of a certain "local gauge symmetry" as was discussed by Hata.

---

[*] Talk given at International Symposium on BRS Symmetry, Sept. 18 – 22, 1995, Kyoto.

[†] E-mail address: kugo@gauge.scphys.kyoto-u.ac.jp

1 Introduction

It is a well-known consequence of the Slavnov-Taylor identity that the ratio of vertex renormalization factor to wave-function renormalization factor is *universal*:

$$\frac{Z_1}{Z_3} = \frac{\widetilde{Z}_1}{\widetilde{Z}_3} = \frac{Z_{\bar{\psi}\psi A}}{Z_\psi} = \ \cdots \ , \tag{1.1}$$

where the denominators Z_3, $\widetilde{Z}_3$ and Z_ψ are the wave-function renormalization factors of gauge-boson, Faddeev-Popov ghost and matter field ψ, respectively, and the numerators Z_1, $\widetilde{Z}_1$ and $Z_{\bar{\psi}\psi A}$ are the gauge-boson vertex renormalization factors of those fields. Namely, the ratio is independent of the measured matter fields (and equals 1 in the QED case as a result of the Ward identity). [We may, however, have to keep in mind that it is *gauge-dependent*.]

The main purpose of this talk is to show that the following equality holds for this universal renormalization factor Z_1/Z_3 generally in covariant gauges (with arbitrary gauge parameter α):

$$\frac{Z_1}{Z_3} = 1 + u \ , \tag{1.2}$$

where $u \equiv u(p^2\!=\!0)$ and $u(p^2)$ is the function defined by

$$\int d^4x \, e^{ipx} \, \langle 0|\, \mathrm{T} D_\mu c^a(x) \, D_\nu \bar{c}^b(0) \,|0\rangle = \left[\Big(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \Big) u(p^2) - \frac{p_\mu p_\nu}{p^2} \right] \delta_{ab} \ . \tag{1.3}$$

This relation (1.2) is very interesting since it is known[1] that $1+u=0$ gives a sufficient condition for all the colored states to become unphysical; namely,

$$1 + u = 0 \quad \Longrightarrow \quad \text{Color Confinement}\,! \tag{1.4}$$

As a preparation for it, we briefly explain in Sect. 2 why the condition $1+u=0$ implies the color confinement. And then in Sect. 3 we give a proof of the relation (1.2). Some implications are discussed in Sect. 4, where I give an argument that the confinement condition $1+u=0$ implies the disappearance of the massless (vector) gauge boson pole. It, therefore, turns to imply that a certain "local gauge symmetry" is dynamically restored as was discussed by Hata[7]. This is explained in Sect. 5. The final Sect. 6 is devoted to some further discussions.

2 $1+u=0$: A Color Confinement Condition

Let us first recapitulate how the condition $1+u=0$ is related to the color confinement[1]. As noted by Ojima[2] first, the equation of motion for the gauge field can be written in the following form of Maxwell-type:

$$gJ^a_\mu = \partial^\nu F^a_{\mu\nu} + \{Q_\mathrm{B}, \, D_\mu \bar{c}^a\} \ , \tag{2.1}$$

where J^a_μ is the Noether current of color symmetry (global gauge symmetry). The point is that the Noether current always has an *arbitrariness* adding a term of the form $\partial^\nu f_{[\mu\nu]}$ with an arbitrary local anti-symmetric tensor $f_{[\mu\nu]}$. That is, the modified current $J'^a_\mu \equiv J^a_\mu + \partial^\nu f_{[\mu\nu]}$ is still conserved and, moreover, the corresponding charge generates the infinitesimal color rotation correctly on any field operators (at least with formal application of canonical commutation relations). We have, therefore, a possibility to define the color charge Q^a by

$$\begin{aligned} Q^a &= \int d^3x \left(J^a_\mu - \frac{1}{g}\partial^\nu F^a_{\mu\nu} \right)_{\mu=0} \\ &= \int d^3x \, \frac{1}{g}\{Q_\mathrm{B},\, D_\mu \bar{c}^a\}_{\mu=0} \end{aligned} \tag{2.2}$$

If we *could* define the color charge by this equation, then it takes a BRS-exact form and so the color confinement is concluded: indeed, for any physical states $|\mathrm{phy}\rangle$ specified by the condition $Q_\mathrm{B}\,|\mathrm{phys}\rangle = 0$, we have

$$\langle \mathrm{phys}|\, Q^a \,|\mathrm{phys}'\rangle = 0 \ . \tag{2.3}$$

It is an easy exercise to show from this equation that *all the physical (that is, BRS-singlet) particles are color-singlet*[1].

The expression Eq. (2.2), however, does *not* give a well-defined color charge operator generally. This is because there is a massless one-particle contribution to J^a_μ, $\partial^\nu F^a_{\mu\nu}$ and $\{Q_\mathrm{B},\, D_\mu \bar{c}^a\}$ so that the 3-volume integration does not converge. To show this, we have first to explain the elementary quartet (a quartet = a pair of BRS-doublets).

We can easily show that there always exists a massless quartet for each group index a. Indeed, using the equation of motion $\partial^\mu D_\mu c^a = 0$ and equal time anti-commutation relation, we find an identity

$$\partial^\mu_x \langle 0|\, \mathrm{T} D_\mu c^a(x)\, \bar{c}^b(y)\, |0\rangle = \delta^{ab}\delta^4(x-y) \ , \tag{2.4}$$

implying that

$$\int d^4x\, e^{ipx} \langle 0|\, \mathrm{T} D_\mu c^a(x)\, \bar{c}^b(0)\, |0\rangle = i\delta^{ab}\frac{p_\mu}{p^2} \ . \tag{2.5}$$

But, an identity $\langle 0|\, \{Q_\mathrm{B}, \mathrm{T}(A^a_\mu\, \bar{c}^b)\}\, |0\rangle = 0$ means the Ward-Takahashi identity

$$i\,\langle 0|\, \mathrm{T} D_\mu c^a(x)\, \bar{c}^b(y)\, |0\rangle = \langle 0|\, \mathrm{T} A^a_\mu(x)\, B^b(y)\, |0\rangle \ , \tag{2.6}$$

so that Eq. (2.5) implies also an identity:

$$\int d^4x\, e^{ipx} \langle 0|\, \mathrm{T} A^a_\mu(x)\, B^b(0)\, |0\rangle = -\delta^{ab}\frac{p_\mu}{p^2} \ . \tag{2.7}$$

The identities (2.5) and (2.7) give exact Green functions and so the presence of massless pole $\propto 1/p^2$ implies that the following four massless asymptotic fields γ^a, $\bar{\gamma}^a$, χ^a and β^a exist for each group index a:

$$\begin{aligned} D_\mu c^a(x) &\xrightarrow[x_0\to\pm\infty]{} \partial_\mu \gamma^a(x) + \cdots \\ \bar{c}^a(x) &\longrightarrow \bar{\gamma}^a(x) + \cdots \\ A^a_\mu(x) &\longrightarrow \partial_\mu \chi^a(x) + \cdots \\ B^a(x) &\longrightarrow \beta^a(x) + \cdots \ . \end{aligned} \tag{2.8}$$

From BRS transformation law $[iQ_{\rm B},\, A^a_\mu] = D_\mu c^a$ and $\{Q_{\rm B},\, \bar{c}^a\} = B^a$, we find that these asymptotic fields transform as follows under the BRS transformation:

$$[iQ_{\rm B},\, \chi^a(x)] = \gamma^a(x) \ , \qquad \{Q_{\rm B},\, \bar{\gamma}^a(x)\} = \beta^a(x) \ . \tag{2.9}$$

This clearly shows that these asymptotic fields γ^a, $\bar{\gamma}^a$, χ^a and β^a really forms a BRS quartet representation.

Now we come back to the well-definedness problem of the expression Eq. (2.2). The point is that $D_\mu \bar{c}^a = \partial_\mu \bar{c}^a + g(A_\mu \times \bar{c})^a$ generally contains the one-particle contribution of the massless asymptotic field $\bar{\gamma}^a$:

$$D_\mu \bar{c}^a = \partial_\mu \bar{c}^a + g(A_\mu \times \bar{c})^a \quad \longrightarrow \quad (1+u)\partial_\mu \bar{\gamma}^a \ , \tag{2.10}$$

where the weight 1 comes from the first term $\partial_\mu \bar{c}^a$ and the weight u from the second $g(A_\mu \times \bar{c})^a$. This can be seen by looking at the definition (1.3) of u and Eq. (2.5). We thus see that the operator $\{Q_{\rm B},\, D_\mu \bar{c}^a\}$ contains the massless one-particle contribution of the elementary quartet member β^a in the form

$$\{\, Q_{\rm B},\, D_\mu \bar{c}^a(x)\,\} \quad \xrightarrow[x_0 \to \pm\infty]{} \quad (1+u)\partial_\mu \beta^a(x) \ . \tag{2.11}$$

Generally, the other operators J^a_μ and $\partial^\nu F^a_{\mu\nu}$ appearing in the Maxwell equation (2.1) also have the one-particle contributions from β^a since they carry the same quantum numbers as $\{Q_{\rm B},\, D_\mu \bar{c}^a\}$:

$$J^a_\mu(x) \quad \longrightarrow \quad v\partial_\mu \beta^a(x) \ , \qquad \partial^\nu F^a_{\mu\nu}(x) \quad \longrightarrow \quad -w\partial_\mu \beta^a(x) \ , \tag{2.12}$$

with suitable weights v and w. In QED, one can easily see that $u = 0$ and $w = Z_3$. The Maxwell equation (2.1) tells us the relation between those weights:

$$gv = -w + (1+u) \ . \tag{2.13}$$

The well-defined color charge is, therefore, given by

$$Q^a = \int d^3x \left(J^a_\mu + \frac{v}{w} \partial^\nu F^a_{\mu\nu} \right)_{\mu=0} \ , \tag{2.14}$$

so that the massless one-particle contribution $\partial_\mu \beta^a$ contained in J^a_μ is cancelled by that in $\partial^\nu F^a_{\mu\nu}$.

Now, if the condition $1 + u = 0$ holds, then we have $v/w = -1/g$ and so the well-defined color charge Eq. (2.14) happens to coincide with the previous BRS-exact expression (2.2) implying the color confinement.

3 Proof of the relation $Z_1/Z_3 = 1 + u$

I have noticed this relation $Z_1/Z_3 = 1 + u$ for the first time in studying the renormalization problem in the background field method in collaboration with Imamura and Van Proeyen[3].



Figure 1: Diagram of massless pole contribution.

However, the proof by that method is a bit complicated and so here I will give a simpler proof using a similar method to Aoki's[4] which he used when proving the electro-magnetic charge universality in the Weinberg-Salam model.

Add $g(v/w)\partial^\nu F^a_{\mu\nu}$ to both sides of the Maxwell equation (2.1), and then we have

$$gJ^{a,\text{well-def}}_\mu = \Big(1+g\frac{v}{w}\Big)\partial^\nu F^a_{\mu\nu} + \{Q_{\mathrm{B}},\, D_\mu \bar{c}^a\}\ , \tag{3.1}$$

with

$$J^{a,\text{well-def}}_\mu \equiv J^a_\mu + \frac{v}{w}\partial^\nu F^a_{\mu\nu} \tag{3.2}$$

being the color current giving the well-defined color charge (2.14). Using a relation $1+gv/w = (1+u)/w$ following from Eq. (2.13), we find

$$\frac{1+u}{w}\partial^\nu F^a_{\mu\nu} = gJ^{a,\text{well-def}}_\mu - \{Q_{\mathrm{B}},\, D_\mu \bar{c}^a\}\ . \tag{3.3}$$

Sandwitching this with two physical states $\langle f|$ and $|g\rangle$ satisfying $\langle f|\, Q_{\mathrm{B}} = Q_{\mathrm{B}}\, |g\rangle = 0$, yields

$$\frac{1+u}{w}\partial^\nu \langle f|\, F^a_{\mu\nu}(x)\, |g\rangle = g\, \langle f|\, J^{a,\text{well-def}}_\mu(x)\, |g\rangle\ . \tag{3.4}$$

We now make an operation $\lim_{p\to 0}\int d^4x\, e^{ipx}$ act on both sides of this equation and evaluate the both sides separately. Start with the left-hand side. Because of the presence of the divergence ∂^ν, only massless one-particle pole contribution to the matrix element $\langle f|\, F^a_{\mu\nu}(x)\, |g\rangle$ can survive in this limit. But such a massless one-particle pole (if any) comes from the gauge boson contribution as shown in Fig. 1 and generally takes the form

$$\begin{aligned}
&\int d^4x\, e^{ipx}\, \langle f|\, F^a_{\mu\nu}(x)\, |g\rangle \Big|_{\text{pole part}} = \Big\langle\, F^a_{\mu\nu}\, A^b_{\mathrm{r},\rho}\,\Big\rangle\, V^{\rho,b}_{fi}\ ,\\
&\quad \Big\langle\, F^a_{\mu\nu}\, A^b_{\mathrm{r},\rho}\,\Big\rangle = -Y\frac{p_\mu g_{\nu\rho} - p_\nu g_{\mu\rho}}{p^2}\delta_{ab}\ ,\\
&\quad V^{\rho,b}_{fi} = ig_{\mathrm{r}} T^b_{fi}(p_f+p_i)^\rho (2\pi)^4 \delta^4(p+p_f-p_i)\ ,
\end{aligned} \tag{3.5}$$

where $\Big\langle\, F^a_{\mu\nu}\, A^b_{\mathrm{r},\rho}\,\Big\rangle$ stands for the propagator from $F^a_{\mu\nu}$ to the renormalized gauge boson $A^b_{\mathrm{r},\rho}$, and $V^{\rho,b}_{fi}$ for the vertex of the renormalized gauge boson between the on-shell initial and final states. Y is a weight factor and g_r is the renormalized coupling constant. Substituting this expression, we find that the left-hand side becomes

$$-Y\frac{1+u}{w}\lim_{p\to 0}\Big(-i\frac{p_\mu p_\rho}{p^2}+ig_{\mu\rho}\Big)(ig_\mathrm{r})T^a_{fi}(p_f+p_i)^\rho(2\pi)^4\delta^4(p+p_f-p_i)\ . \tag{3.6}$$

Noting the pole term $p_\mu p_\rho/p^2$ vanishes owing to the "current conservation" at the vertex, we obtain

$$\begin{aligned}\frac{1+u}{w}\lim_{p\to 0}\int d^4x\, e^{ipx}\partial^\nu\,\langle f|\, F^a_{\mu\nu}(x)\,|g\rangle& \\ =\ Y\frac{1+u}{w}g_\mathrm{r}T^a_{fi}\lim_{p_f\to p_i}(p_f+p_i)_\mu(2\pi)^4\delta^4(p_f-p_i)\ .&\end{aligned} \tag{3.7}$$

The weight factor Y introduced in Eq. (3.5) can easily be related to the previous weight w: using $\partial^\nu F^a_{\mu\nu}(x)\longrightarrow -w\partial_\mu\beta^a(x)$ and the WT identity (2.7) with $A^b_{\mathrm{r},\rho}=Z_3^{-1/2}A^b_\rho$, we obtain

$$\Big\langle\,\partial^\nu F^a_{\mu\nu}\,A^b_{\mathrm{r},\rho}\,\Big\rangle\Big|_{\text{pole part}}=-w\,\Big\langle\,\partial_\mu B^a\,A^b_{\mathrm{r},\rho}\,\Big\rangle=wZ_3^{-1/2}(i\delta^{ab})\frac{p_\mu p_\rho}{p^2}\ , \tag{3.8}$$

comparison of which with the divergence of the second equation in Eq. (3.5) leads to

$$Y=wZ_3^{-1/2}\ ,\qquad \text{so that}\qquad Y\frac{1+u}{w}=(1+u)Z_3^{-1/2}. \tag{3.9}$$

The right hand side of Eq. (3.4), on the other hand, can be easily evaluated for $\mu=0$ component as follows:

$$\begin{aligned}\lim_{p\to 0}\int d^4x\, e^{ipx}g\,\langle f|\,J_0^{a,\text{well-def}}(x)\,|g\rangle &= \int dx_0\, g\,\langle f|\,Q^{a,\text{well-def}}\,|g\rangle \\ &= gT^a_{fi}\lim_{p_f\to p_i}2p_i^0(2\pi)^4\delta^4(p_f-p_i)\ .\end{aligned} \tag{3.10}$$

Now comparing this with the above Eq. (3.7) with Eq. (3.9), we finally obtain a relation

$$(1+u)Z_3^{-1/2}g_\mathrm{r}=g\ . \tag{3.11}$$

But, if we use the relation $g=Z_1Z_3^{-3/2}g_\mathrm{r}$ between the bare and renormalized coupling constants g and g_r, this is nothing but the desired relation $1+u=Z_1/Z_3$, Eq. (1.2).

We have proved the relation $1+u=Z_1/Z_3$ in covariant gauges with arbitrary values of gauge parameter α. In Landau gauge $\alpha=0$, however, it is easier to show it. First define the following two-point functions

$$\begin{aligned}\langle\, c\;\bar c\,\rangle\ &\equiv\ -\frac{1}{p^2G(p^2)}\ ,\\ \langle\, g(A_\mu\times c)\;\bar c\,\rangle_{\text{1PI}}\ &\equiv\ \langle\, g(A_\mu\times c)\;\bar c\,\rangle\,/\,\langle\, c\;\bar c\,\rangle\equiv -ip_\mu F(p^2)\ .\end{aligned} \tag{3.12}$$

Here and henceforth we use an abbreviated notation:

$$\langle\ A\,B\ \rangle\equiv\int d^4x\, e^{ipx}\,\langle 0|\,\mathrm{T}\,A(x)\,B(0)\,|0\rangle\ . \tag{3.13}$$

The suffix 1PI generally denotes the one-particle irreducible vertex. Then, we have

$$\langle\, D_\mu c\ \bar{c}\,\rangle = \langle\, \partial_\mu c\ \bar{c}\,\rangle + \langle\, g(A_\mu\times c)\ \bar{c}\,\rangle \equiv ip_\mu(1+F(p^2))\frac{1}{p^2G(p^2)}\ . \tag{3.14}$$

But this must equal ip^μ/p^2 because of Eq. (2.7) so that

$$1+F(p^2)=G(p^2)\ . \tag{3.15}$$

On the other hand, a simple diagramatical consideration shows that

$$\langle\, c\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI} = ip^\rho\,\langle\, g(A_\rho\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI} \tag{3.16}$$

and hence that

$$\begin{aligned}\langle\, D_\mu c\ g(A_\nu\times\bar{c})\,\rangle &= \langle\, D_\mu c\ \bar{c}\,\rangle\ \langle\, c\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI} + \langle\, g(A_\mu\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI} \\ &= \Big(\delta_\mu^\rho - \frac{p_\mu p^\rho}{p^2}\Big)\,\langle\, g(A_\rho\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI}\ .\end{aligned} \tag{3.17}$$

The last Green function $\langle\, D_\mu c\ g(A_\nu\times\bar{c})\,\rangle$ must be equal to $(g_{\mu\nu}-(p_\mu p_\nu/p^2))u(p^2)$ because of the definition (1.3) of the function $u(p^2)$ so that $\langle\, g(A_\mu\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI}$ have to have the form

$$\langle\, g(A_\mu\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI} = g_{\mu\nu}u(p^2)+p_\mu p_\nu v(p^2) \tag{3.18}$$

with $v(p^2)$ being an arbitrary function. Up to here all the equations hold for any covariant α gauges. But, in Landau gauge $\alpha=0$, we have additionally an identity

$$\begin{aligned}-\,ip_\mu F(p^2) &\equiv \langle\, g(A_\mu\times c)\ \bar{c}\,\rangle_{\rm 1PI} \\ &\equiv \langle\, g(A_\mu\times c)\ g(A_\nu\times\bar{c})\,\rangle_{\rm 1PI}\times(-ip^\nu)\ .\end{aligned} \tag{3.19}$$

This is because the derivative factor acting on the anti-ghost at the $\bar{c}$-A_ν-c vertex at the right end of the diagrams can be transferred to act on the external ghost since $\partial_\nu A^\nu=0$ in the Landau gauge. Therefore we have

$$-\,ip_\mu F(p^2) = -ip_\mu\Big(u(p^2)+p^2v(p^2)\Big)\ ,\quad \Longrightarrow\quad G(p^2)-1=u(p^2)+p^2v(p^2)\ . \tag{3.20}$$

This relation yields at $p^2=0$

$$G(0)=1+u(0)\ , \tag{3.21}$$

but, since $G(0)=\widetilde{Z}_3^{-1}$ and $\widetilde{Z}_1=1$ in the Landau gauge, this shows the relation $\widetilde{Z}_1/\widetilde{Z}_3=1+u$, thus proving the desired identity Eq. (1.2).

4 What Does $1+u=Z_1/Z_3$ Imply?

How is $1+u=Z_1/Z_3=0$ expected to be realized? Naively, we immediately expect the following two possibilities:

1. $Z_1=0$, and $Z_3=$ finite.

2. $Z_1 =$ finite, and $Z_3 = 0$.

However, since the condition is concerned with only the ratio, we have, for instance, even the possibility

3. $Z_3 = 0$, but/and $Z_1/Z_3 = 0$.

Here I mean that Z_1 has a higher zero than Z_3; *e.g.*, $Z_3(p^2) = \mathcal{O}(p^2)$ and $Z_1(p^2) = \mathcal{O}(p^4)$. The wave-function renormalization factor $Z_3(p^2)$ as a function of p^2 can be defined as follows by the general form of gauge boson 2-point function:

$$\int d^4x\, e^{ipx} \langle 0|\, \mathrm{T}\, A^a_\mu(x)\, A^b_\nu(x)\, |0\rangle = \left[\Big(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \Big) \frac{Z_3(p^2)}{p^2} + \alpha \frac{p_\mu p_\nu}{p^4} \right] \delta^{ab} \tag{4.1}$$

Although I cannot prove it, it seems plausible that the possibility 3 is realized when $1+u = 0$. Let us now explain this.

Assume that $Z_3(p^2=0) \neq 0$, then, there exists a *vector* asymptotic field $\mathcal{A}^a_\mu(x)$ in the channel $A^a_\mu(x)$:

$$A^a_\mu(x) \quad \xrightarrow[x_0 \to \pm\infty]{} \quad \mathcal{A}^a_\mu(x)\ . \tag{4.2}$$

This is a colored particle. But any colored particles are confined when $1+u = 0$. Therefore the BRS transform of $A^a_\mu(x)$

$$[iQ_{\mathrm{B}},\, A^a_\mu(x)] = D_\mu c^a(x)\ , \tag{4.3}$$

with ghost number 1, should also have its own massless vector asymptotic field $\mathcal{C}^a_\mu(x)$ such that $(\mathcal{A}^a_\mu(x),\, \mathcal{C}^a_\mu(x))$ form a BRS-doublet (and hence become unphysical):

$$[iQ_{\mathrm{B}},\, \mathcal{A}^a_\mu(x)] = \mathcal{C}^a_\mu(x)\ . \tag{4.4}$$

This is a color confinement by the quartet mechanism[5]. Since this asymptotic field $\mathcal{C}^a_\mu(x)$ is a vector, it must be a bound-state in the $(A_\mu \times c)^a$ channel. So it is natural* to suppose that it should show up as a pole in the 1PI Green function $\langle\, g(A_\mu{\times}c)\ g(A_\nu{\times}\bar{c})\, \rangle_{\mathrm{1PI}}$, namely, in Eq. (3.18)

$$\langle\, g(A_\mu{\times}c)\ g(A_\nu{\times}\bar{c})\, \rangle_{\mathrm{1PI}} = g_{\mu\nu} u(p^2) + p_\mu p_\nu v(p^2)\ .$$

The vector massless pole $1/p^2$ should appear in the $g_{\mu\nu}$-part, *i.e.*, in $u(p^2)$. But this contradicts with the present assumption $u = u(p^2{=}0) = 1$.

This argument, therefore, suggests that the massless vector pole in A^a_μ, which existed in the perturbation phase, should disappear (or the appearance of mass gap): namely,

$$Z_3(p^2) \propto p^2\ , \tag{4.5}$$

corresponding to the above possibility 3.

---

[*] This is the point which is not quite rigorous in this argument. There remains a possibility that the vector asymptotic field $\mathcal{C}^a_\mu(x)$ exists but nevertheless does not produce a pole in the 1PI Green function $\langle\, g(A_\mu{\times}c)\ g(A_\nu{\times}\bar{c})\, \rangle_{\mathrm{1PI}}$. This might not be so strange a possibility since we are now considering in any case such an unfamiliar situation that the always existing quartet member β^a disappears from the operator $\{Q_{\mathrm{B}},\, D_\mu \bar{c}^a\}$. This point was emphasized by Izawa[6]. But, if $\langle\, g(A_\mu{\times}c)\ g(A_\nu{\times}\bar{c})\, \rangle_{\mathrm{1PI}}$ does not show up the massless vector pole, then, where will the asymptotic vector field $\mathcal{C}^a_\mu(x)$ produce its pole at all?

5 Dynamical Restoration of a "Local" Gauge Symmetry

The conclusion in the previous section reminds me Hata's work[7, 8] who clarified a symmetry aspect of the 'strange' condition $1+u=0$. So let us briefly recapitulate what he has done in this context.

In QED, the covariant α gauge fixing still leaves a symmetry under the gauge transformation with transformation parameter linear in x:

$$\Lambda(x) = a_\nu x^\nu + b \ . \tag{5.1}$$

Although being a (special) "local" gauge symmetry, this transformation can be regarded as a combination of global transformations with x-independent parameters a_ν and b. We have accordingly conserved Noether charges, vector one Q^ν and scalar one Q, (the latter Q being just a usual electro-magnetic (global $U(1)$) charge), and they give the generators of the original transformation:

$$\begin{aligned} &[\, i(a_\nu Q^\nu + bQ),\, A_\mu(x)\,] = \partial_\mu \Lambda(x) = a_\mu \ , \\ &\Longrightarrow \quad \begin{cases} [\, iQ,\, A_\mu(x)\,] = 0 \\ [\, iQ^\nu,\, A_\mu(x)\,] = \delta^\nu_\mu \end{cases} \end{aligned} \tag{5.2}$$

The first commutation relation says only that the photon A_μ carries vanishing charge, but the second one shows that the vector symmetry of charge Q^ν is *always* spontaneously broken since the right-hand side is non-vanishing c-number δ^ν_μ! It also says that the massless Nambu-Goldstone (NG) mode corresponding to the spontaneous breaking should appear in the A_μ channel. It can be argued that, as far as the global $U(1)$ symmetry corresponding to the scalar charge Q is not spontaneously broken, the NG mode must be a vector particle so that the photon can be identified with the Nambu-Goldstone vector[9]! [This type of argument is important since it *proves* the existence and exact masslessness of the photon. Quite a parallel argument can apply to gravity[10, 11] and proves the existence and exact masslessness of the graviton.]

Hata considered the same thing in non-Abelian gauge theory. Corresponding to the "local" gauge transformation,

$$\Lambda^a(x) = a^a_\nu x^\nu + b^a \ , \tag{5.3}$$

there appear the following scalar and vector charges, both of which carry color index a now:

$$\begin{aligned} Q^a \quad &\longrightarrow \quad J^a_\mu(x) \ , \\ Q^{a,\nu} \quad &\longrightarrow \quad J^{a,\nu}_\mu(x) = gJ^a_\mu(x)x^\nu + F^{a\ \nu}_{\mu} \ . \end{aligned} \tag{5.4}$$

In this case, however, the vector transformation with parameter a^a_ν is not an exact symmetry but is slightly violated and so

$$\partial^\mu J^a_{\mu,\nu}(x) = \{Q_\mathrm{B},\, D_\nu \bar{c}^a\} \ . \tag{5.5}$$

In his second paper[8], Hata considered a modified vector transformation which becomes an exact symmetry. But here we continue to consider the present simplified version. Again, this "local symmetry" with parameter $a^a_\nu x^\nu$ is spontaneously broken in the perturbation phase and the usual perturb-ative massless gauge boson can be identified with Nambu-Goldstone vector.

Contrary to the QED case, however, there is a possibility that this "local symmetry" is restored dynamically in the non-Abelian case. Hata's picture for the color confinement is as follows:

$$\begin{aligned}&\text{Dynamical restoration of the "local symmetry" with parameter } a^a_\nu x^\nu \\ &= \quad \text{Disordered phase} \quad = \quad \text{Confinement phase} .\end{aligned} \tag{5.6}$$

Indeed, if the "local symmetry" is dynamically restored, then the massless vector pole (*i.e.*, perturbative gauge boson pole) in the current $J^a_{\mu,\nu}$ should disappear and it will have no massless pole contribution. This requirement actually leads to the above condition $1+u=0$ of ours, since we have

$$\partial^\mu \left\langle \; J^a_{\mu,\nu}(x) \; A^b_\rho \; \right\rangle = i\Big(g_{\mu\rho} - \frac{p_\mu p_\rho}{p^2}\Big)(1+u(p^2)) \; . \tag{5.7}$$

Namely our condition $1+u=0$ is a necessary condition for the restoration of the "local symmetry". I do not know whether it is also sufficient, but our discussion in the previous section strongly suggests that it *is*. Hata also gave another interesting direct proof that any colored particle has to be a BRS-doublet if the current $J^a_{\mu,\nu}(x)$ has no massless pole, by considering a 3-point Green function of $J^a_{\mu,\nu}(x)$ and the two fields corresponding to that colored particle[7].

6 Discussions

One may wonder how it is possible at all to satisfy our confinement condition $1+u=0$. This question is quite natural if we note the fact that $u = u(p^2=0)$ is both infrared and ultraviolet divergent in perturbation theory!

However, we should recall that quite a similar thing is actually realized in the non-linear sigma models in two dimension. As an example, consider the following $O(N)/O(N-1)$ sigma model again following Hata[7]:

$$\mathcal{L} = \frac{1}{2}\frac{(\partial\boldsymbol{\phi})^2}{\Big(1+\dfrac{\lambda}{4}\boldsymbol{\phi}^2\Big)^2} \; , \qquad\qquad \boldsymbol{\phi} \equiv \begin{pmatrix} \phi_1 \\ \vdots \\ \phi_{N-1} \end{pmatrix} \; . \tag{6.1}$$

The non-linearly realized $O(N)/O(N-1)$ symmetry is the transformation:

$$\delta_i \phi_j = \Big(1 - \frac{\lambda}{4}\boldsymbol{\phi}^2\Big)\delta_{ij} + \frac{\lambda}{2}\phi_i\phi_j \; , \tag{6.2}$$

with $i = 1,\ 2,\ \cdots,\ N-1$. In perturbation phase, this non-linear symmetry is (spontaneously) broken and the fields ϕ_i are the corresponding massless NG bosons. The restoration condition for this symmetry is given by, in the leading order in $1/N$ expansion,

$$\langle\, \delta_i \phi_j \,\rangle = \delta_{ij}\Big(1 - \frac{\lambda}{4}\left\langle \boldsymbol{\phi}^2 \right\rangle\Big) = 0 \ . \tag{6.3}$$

This is quite analogous to our condition $1+u=0$. The vacuum expectation value $-(\lambda/4)\left\langle \boldsymbol{\phi}^2 \right\rangle$ corresponds to our u, and is indeed both infrared and ultraviolet divergent in perturbation calculation! Nevertheless we know[12] that $(1-\lambda\left\langle \boldsymbol{\phi}^2 \right\rangle/4)$ is actually realized in this model. It is known rather generally that a mass gap appears and the nonlinear symmetry is dynamically restored in a wide variety of two dimensional non-linear sigma models.

As a matter of fact, this resemblance of dynamical symmetry restorations between Yang-Mills gauge theory in four dimension and non-linear sigma model in two dimension, is not a mere analogy. Indeed, it becomes an exact correspondence if we consider a pure gauge model of Yang-Mills theory in four dimension[13]. Let us close this talk by briefly explaining this.

Take the $SU(N)$ Yang-Mills theory with $OSp(4|2)$ symmetric gauge fixing:

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^2 + \delta_{\rm B}\bar{\delta}_{\rm B}\Big(\frac{1}{2}A_\mu^2 + i\bar{c}c\Big) \tag{6.4}$$

where $\delta_{\rm B}$ and $\bar{\delta}_{\rm B}$ denote BRS and anti-BRS transformations, respectively. The pure gauge model is given by replacing the gauge field A_μ^a in this Lagrangian by the pure gauge mode $g^\dagger\partial_\mu g$:

$$A_\mu(x) \qquad \longrightarrow \qquad g^\dagger(x)\partial_\mu g(x) \ . \tag{6.5}$$

Then, the $F_{\mu\nu}^2$ term drops out and only the gauge-fixing term remains. Namely the pure gauge model is a kind of topological model. One can show very easily that this model is *exactly* equivalent by the Parisi-Sourlas mechanism to the two dimensional $SU(N)_{\rm L} \times SU(N)_{\rm R}/SU(N)$ non-linear sigma model with the action

$$\int d^2x\, {\rm tr}(\partial^\mu g^\dagger \partial_\mu g) \ . \tag{6.6}$$

This model has an $SU(N)_{\rm L} \times SU(N)_{\rm R}$ symmetry under the transformation $g \to h_{\rm L}^\dagger g h_{\rm R}$. Very interestingly, the current corresponding to the $SU(N)_{\rm R}$ symmetry is found to be given by

$$J_{{\rm R}\,\mu}^a = \{Q_{\rm B},\, D_\mu \bar{c}^a\} \ . \tag{6.7}$$

So, if the symmetry is restored, then it implies that our condition $1+u=0$ is literally realized. This is what actually occurs as we know from the exact result due to Polyakov and Wiegmann[14]. So this pure gauge model has a possibility to be used as a “zeroth order” theory in a new form of “perturbation theory” in the actual non-Abelian gauge theory, in which the confinement condition $1+u=0$ is satisfied order by order. It is encouraging that such trials towards this direction are already performed by several authors[15, 16, 17].

Acknowledgements

I would like to thank H. Hata, K. Izawa and Y. Taniguchi for valuable discussions and comments. He is supported in part by the Grant-in-Aid for Cooperative Research (#07304029) and the Grant-in-Aid for Scientific Research (#06640387) from the Ministry of Education, Science and Culture.

References

[1] T. Kugo and I. Ojima, *Prog. Theor. Phys.* Supplement **66** (1979) 1.

[2] I. Ojima, *Nucl. Phys.* **B143** (1978) 340.

[3] Y. Imamura, T. Kugo and A. Van Proeyen, "S–Matrix from Effective Action in Background Field Method", Kyoto preprint in preparation.

[4] K.I. Aoki, *Soryusiron Kenkyu* (in Japanese) **59** (1979) 383.

[5] T. Kugo, *Phys. Lett.* **83B** (1979) 93.

[6] K.I. Izawa, private communication after my talk.

[7] H. Hata, *Prog. Theor. Phys.* **67** (1982) 1607.

[8] H. Hata, *Prog. Theor. Phys.* **69** (1983) 1524.

[9] R. Ferrari and L.E. Picasso, *Nucl. Phys.* **B31** (1971) 316.
R.A. Brandt and Ng Wing-Chiu, *Phys. Rev.* **D10** (1974) 4198.

[10] N. Nakanishi and I. Ojima, *Phys. Rev. Lett.* **43** (1979) 91.

[11] T. Kugo, H. Terao and S. Uehara, *Prog. Theor. Phys.* Supplement **85** (1985) 122.

[12] W.A. Bardeen, B.W. Lee and R.E. Shrock, *Phys. Rev.* **D14** (1976) 985.

[13] H. Hata and T. Kugo, *Phys. Rev.* **D32** (1985) 938.

[14] A. Polyakov and P.B. Wiegmann, *Phys. Lett.* **131B** (1983) 121.
P.B. Wiegmann, *Phys. Lett.* **141B** (1984) 217; **142B** (1984) 173.

[15] H. Abe and N. Nakanishi, *Prog. Theor. Phys.* **89** (1993) 501.

[16] K.I. Izawa, *Prog. Theor. Phys.* **90** (1993) 911.

[17] H. Hata and Y. Taniguchi, *Prog. Theor. Phys.* **94** (1995) 435.